

**CAMBRIDGE INTERNATIONAL EXAMINATIONS**  
GCE Advanced Subsidiary Level and GCE Advanced Level

## **MARK SCHEME for the May/June 2014 series**

### **9701 CHEMISTRY**

**9701/21**

Paper 2 (Structured Questions AS Core),  
maximum raw mark 60

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2014 series for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level components and some Ordinary Level components.

|        |                                |          |       |
|--------|--------------------------------|----------|-------|
| Page 2 | Mark Scheme                    | Syllabus | Paper |
|        | GCE AS/A LEVEL – May/June 2014 | 9701     | 21    |

| Question | Mark Scheme  | Mark        | Total |
|----------|--|-------------|-------|
| 1 (a)    | The amount of energy required / energy change / enthalpy change when one electron is removed<br>from each atom / (cat)ion in one mol<br>of gaseous atoms / (cat)ions<br><b>OR</b> energy change when 1 mole of electrons is removed from one mole of gaseous atoms / ions<br>$X(g) \rightarrow X^+(g) + e^-$ gains 2 marks | 1<br>1<br>1 | 3     |
| (b) (i)  | Group V / 5 / 15<br><br>Big difference between fifth and sixth ionisation energies   | 1<br>1      | 2     |
| (ii)     | $1s^2 2s^2 2p^3$<br>ecf from (b)(i) if period 2  | 1           | 1     |
| (c) (i)  | (Weighted) mean / average mass of an atom(s) (of an element)<br><br>Relative to $1/12^{\text{th}}$ of (the mass of an atom of) carbon-12 <b>OR</b><br>relative to carbon-12 which is (exactly) 12 (units)<br>allow as an expression  | 1<br>1      | 2     |
| (ii)     | $\frac{Z}{A_r} \quad \frac{Cl}{35.5} = 1:2$<br>$\text{So } \frac{68.87/35.5}{31.13/A_r} = 2$<br>$A_r = \frac{2 \times 31.13 \times 35.5}{68.87} = 32.0923 = 32.1 \text{ to } 3\text{s.f.}$<br>Allow alternative correct methods  | 1<br>1      | 2     |

|        |                                |          |       |
|--------|--------------------------------|----------|-------|
| Page 3 | Mark Scheme                    | Syllabus | Paper |
|        | GCE AS/A LEVEL – May/June 2014 | 9701     | 21    |

| Question  | Mark Scheme   | Mark             | Total     |
|-----------|---|------------------|-----------|
| (d) (i)   | $\text{NaCl} (+ \text{aq}) \rightarrow \text{Na}^+ + \text{Cl}^-$<br>$\text{NaCl} + \text{H}_2\text{O} \rightarrow \text{Na}^+ + \text{Cl}^- + \text{H}_2\text{O}$<br><br>$\text{SiCl}_4 + 2\text{H}_2\text{O} \rightarrow \text{SiO}_2 + 4\text{HCl}$<br>$\text{SiCl}_4 + 4\text{H}_2\text{O} \rightarrow \text{Si(OH)}_4 + 4\text{HCl}$<br>$\text{SiCl}_4 + 4\text{H}_2\text{O} \rightarrow \text{SiO}_2 \cdot 2\text{H}_2\text{O} + 4\text{HCl}$<br><br>Allow correct equation with other molar amounts of water | 1<br><br>1       | 2         |
| (ii)      | NaCl is ionic <b>AND</b> giant/lattice<br>NaCl dissolves/does not react<br>SiCl <sub>4</sub> is <u>covalent</u> <b>AND</b> molecular/simple<br>SiCl <sub>4</sub> is hydrolysed/reacts   | 1<br>1<br>1<br>1 | 4         |
| (e)       | shape of SF <sub>6</sub> = Octahedral<br>bond angle = 90°   | 1<br>1           | 2         |
|           |   |                  | <b>18</b> |
| 2 (a) (i) | (The MnO <sub>4</sub> <sup>-</sup> ions cause the Fe <sup>2+</sup> ions to) lose electrons owtte/ora  | 1                | 1         |
| (ii)      | $\text{MnO}_4^-(\text{aq}) + 5\text{Fe}^{2+}(\text{aq}) + 8\text{H}^+(\text{aq}) \rightarrow \text{Mn}^{2+}(\text{aq}) + 5\text{Fe}^{3+}(\text{aq}) + 4\text{H}_2\text{O}(\text{l})$  | 1+1+1            | 3         |
| (b) (i)   | $\frac{20.0 \times 0.020}{1000} = 4(.00) \times 10^{-4} \text{ (mol)}$  | 1                | 1         |
| (ii)      | MnO <sub>4</sub> <sup>-</sup> : Fe <sup>2+</sup> = 1 : 5<br>so amount of Fe <sup>2+</sup> = 5 × 4.00 × 10 <sup>-4</sup> = 2(.00) × 10 <sup>-3</sup> (mol)<br>ecf from (b)(i)  | 1                | 1         |
| (iii)     | $2.00 \times 10^{-3} \times 250/25 = 0.02(00) \text{ (mol)}$<br>ecf from (b)(ii)  | 1                | 1         |

|        |                                |          |       |
|--------|--------------------------------|----------|-------|
| Page 4 | Mark Scheme                    | Syllabus | Paper |
|        | GCE AS/A LEVEL – May/June 2014 | 9701     | 21    |

| Question  | Mark Scheme   | Mark        | Total    |
|-----------|---|-------------|----------|
| (iv)      | $3.40/0.02 = 170$<br>ecf from (b)(iii)  | 1           | 1        |
| (v)       | $170 - 151.8 = 18.2$<br>$18.2/18 = 1.01$<br>$x = 1$<br>ecf from (b)(iv) if appropriate  | 1           | 1        |
|           |   |             | <b>9</b> |
| 3 (a) (i) | <b>K</b> = $Cl^-$ / chloride / $F^-$ / fluoride<br><br>$H_2SO_4 + 2NaCl \rightarrow Na_2SO_4 + 2HCl$ (or equation with F or <b>K</b> for Cl) <b>OR</b><br>$H_2SO_4 + NaCl \rightarrow NaHSO_4 + HCl$ (or equation with F or <b>K</b> for Cl)<br><br>ecf from identity of <b>K</b> so long as halide<br><b>HK</b> is acidic / <b>HK</b> is a gas / an acidic gas is produced | 1<br>1<br>1 | 3        |
| (ii)      | <b>L</b> = $I^-$ / iodide<br><br>colour = yellow<br>ecf from identity of <b>L</b> i.e. $Cl^-$ (white) or $Br^-$ (cream)<br>$Ag^+ + I^- \rightarrow AgI$ (or equation with <b>L</b> )<br>$AgNO_3 + NaI \rightarrow AgI + NaNO_3$ (or equation with <b>L</b> )<br>ecf from identity of <b>L</b> so long as halide   | 1<br>1<br>1 | 3        |
| (iii)     | <u>Br<sub>2</sub></u> / bromine has <b>fewer electrons</b> than iodine / <b>more electrons</b> than chlorine<br><b>intermolecular / van der Waals' forces</b> (in Br <sub>2</sub> / <b>M<sub>2</sub></b> ) <b>weaker</b> than in iodine / <b>stronger</b> than in chlorine  | 1<br>1      | 2        |
| (b) (i)   | <b>B</b> = chlorine / $Cl_2$<br><b>C</b> = hydrogen / $H_2$<br><b>D</b> = sodium hydroxide / NaOH   | 1<br>1<br>1 | 3        |

|        |                                |          |       |
|--------|--------------------------------|----------|-------|
| Page 5 | Mark Scheme                    | Syllabus | Paper |
|        | GCE AS/A LEVEL – May/June 2014 | 9701     | 21    |

| Question | Mark Scheme   | Mark   | Total     |
|----------|---|--------|-----------|
| (ii)     | anode: $2Cl^- \rightarrow Cl_2 + 2e^-$<br>cathode: $2H^+ + 2e^- \rightarrow H_2$ <b>OR</b><br>$2H_2O + 2e^- \rightarrow 2OH^- + H_2$  | 1<br>1 | 2         |
|          |   |        | <b>13</b> |
| 4 (a)    | decolourisation with an alkene at room conditions / quickly / easily /<br><b>OR</b> alkane needs higher temp / UV / is slow at room conditions<br><br>double / $\pi$ / pi bond / C=C present in alkenes | 1<br>1 | 2         |
| (b) (i)  | UV light / sunlight / high temperature  | 1      | 1         |
| (ii)     | (Free) radical<br>Substitution  | 1<br>1 | 2         |
| (iii)    | $\bullet C_2H_5 + \bullet C_2H_5 \rightarrow C_4H_{10}$   | 1      | 1         |
| (iv)     | $C_2H_5Br + Br\bullet \rightarrow \bullet C_2H_4Br + HBr$ <b>OR</b><br>$\bullet C_2H_4Br + Br_2 \rightarrow C_2H_4Br_2 + Br\bullet$   | 1      | 1         |
| (c) (i)  | Electrophilic<br>Addition   | 1<br>1 | 2         |

|        |                                |          |       |
|--------|--------------------------------|----------|-------|
| Page 6 | Mark Scheme                    | Syllabus | Paper |
|        | GCE AS/A LEVEL – May/June 2014 | 9701     | 21    |

| Question | Mark Scheme  | Mark   | Total     |
|----------|--|--------|-----------|
| (ii)     | <p>M1: 2 correct curly arrows</p> <p>M2: correct dipole</p> <p>M3: correct intermediate</p> <p>M4: curly arrow from lone pair on Br<sup>-</sup> to C<sup>+</sup></p> |        | 4         |
| (d)      | <p>minimum of three repeat units</p>   | 2      | 2         |
| (e) (i)  | NaOH/KOH<br>ethanolic/ alcoholic <b>AND</b> heat / reflux  | 1<br>1 | 2         |
| (ii)     |  | 1      | 1         |
| (iii)    | <p>Propanenitrile / propanonitrile / propionitrile / ethyl cyanide / cyanoethane</p>   | 1<br>1 | 2         |
|          |  |        | <b>20</b> |