

**CAMBRIDGE INTERNATIONAL EXAMINATIONS**

Cambridge International Advanced Subsidiary and Advanced Level

**MARK SCHEME for the May/June 2015 series****9701 CHEMISTRY****9701/22**Paper 2 (Structured Questions AS Core),  
maximum raw mark 60

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2015 series for most Cambridge IGCSE<sup>®</sup>, Cambridge International A and AS Level components and some Cambridge O Level components.

® IGCSE is the registered trademark of Cambridge International Examinations.


Page 2	Mark Scheme	Syllabus Paper
	Cambridge International AS/A Level – May/June 2015	9701 / 22

Question	Mark Scheme	Mark	Total																					
1 (a)	name of particle	relative mass	relative charge																					
	proton	1	+1	[1]																				
	electron	1/1836	-1	[1]																				
	neutron	1	0	[1]	[3]																			
(b) (i)	Mass of an atom(s) relative to 1/12 <sup>th</sup> (the mass) of (an atom of) carbon-12 <b>OR</b> relative to carbon-12 which is (exactly) 12	[1]		[1]	[2]																			
	(ii)	% of third isotope = 10 $\frac{(24 \times 79) + (26 \times 11.0) + 10x}{100} = 24.3$ 10x = 248 x = 24.8 (3s.f.)	[1] [1]		[1] [3]																			
(c) (i)	anode $2Cl^- \rightarrow Cl_2 + 2e^-$ cathode $Mg^{2+} + 2e^- \rightarrow Mg$	[1] [1]		[1] [1]	[2]																			
	(ii)	<table border="0"> <tr> <td>Mg</td> <td>O</td> <td>H</td> <td>Cl</td> <td></td> </tr> <tr> <td>31.65</td> <td>20.84</td> <td>1.31</td> <td>46.2</td> <td></td> </tr> <tr> <td>24.3</td> <td>16</td> <td>1</td> <td>35.5</td> <td></td> </tr> <tr> <td>1.30</td> <td>1.30</td> <td>1.31</td> <td>1.30</td> <td>= 1:1:1:1</td> </tr> </table> MgOHCl	Mg	O	H	Cl		31.65	20.84	1.31	46.2		24.3	16	1	35.5		1.30	1.30	1.31	1.30	= 1:1:1:1	[1]	
Mg	O	H	Cl																					
31.65	20.84	1.31	46.2																					
24.3	16	1	35.5																					
1.30	1.30	1.31	1.30	= 1:1:1:1																				
(d) (i)	Na <sub>2</sub> O basic/alkaline; Al <sub>2</sub> O <sub>3</sub> amphoteric/acidic and basic; SO <sub>3</sub> acidic Na <sub>2</sub> O (giant) ionic <b>AND</b> SO <sub>3</sub> (simple/molecular) covalent	[1] [1]		[1] [1]	[2]																			
	(ii)	$Na_2O + 2HCl \rightarrow 2NaCl + H_2O$ $Al_2O_3 + 6HCl \rightarrow 2AlCl_3 + 3H_2O$ $Al_2O_3 + 2NaOH + 7H_2O \rightarrow 2NaAl(OH)_4(H_2O)_2$ <b>OR</b> $Al_2O_3 + 2NaOH + 3H_2O \rightarrow 2NaAl(OH)_4$ <b>OR</b> $Al_2O_3 + 2NaOH \rightarrow 2NaAlO_2 + H_2O$ <b>OR</b> $Al_2O_3 + 2OH^- + 7H_2O \rightarrow 2[Al(OH)_4(H_2O)_2]^-$ <b>OR</b> $Al_2O_3 + 2OH^- + 3H_2O \rightarrow 2[Al(OH)_4]^-$ <b>OR</b> $Al_2O_3 + 2OH^- \rightarrow 2AlO_2^- + H_2O$  $SO_3 + NaOH \rightarrow NaHSO_4$ <b>OR</b> $SO_3 + 2NaOH \rightarrow Na_2SO_4 + H_2O$	[1] [1] [1] [1] [1] [1] [1]		[1] [1] [1] [1] [1] [1] [1]	[4]																		

Page 3	Mark Scheme	Syllabus Paper
	Cambridge International AS/A Level – May/June 2015	9701

Question	Mark Scheme	Mark	Total
			[18]
2 (a) (i)	$2\text{PbS} + 3\text{O}_2 \rightarrow 2\text{PbO} + 2\text{SO}_2$ reagents and formulae balancing	[1] [1]	[2]
(ii)	S (is oxidised) $-2$ to $(+)4$ O (is reduced) $0$ to $-2$	[1] [1]	[2]
(b) (i)	$T = 400 - 600^\circ\text{C}$ (chosen as a compromise because) High T increases rate ora High T decreases yield / moves eqm left / makes less $\text{SO}_3$ as forward reaction exothermic ora	[1] [1] [1]	[3]
(ii)	High pressure increases rate as collision frequency increases ora  High pressure moves eqm right / favours forward reaction as more moles on left ora Uneconomic to use high pressures / high yield at low pressure	[1]  [1] [1]	[3]
(c) (i)	Reaction (too) exothermic / acid spray produced	[1]	[1]
(ii)	$\text{SO}_3 + \text{H}_2\text{SO}_4 \rightarrow \text{H}_2\text{S}_2\text{O}_7$ $\text{H}_2\text{S}_2\text{O}_7 + \text{H}_2\text{O} \rightarrow 2\text{H}_2\text{SO}_4$	[1] [1]	[2]
(d)	Preservative owtte antimicrobial / antioxidant / reducing agent	[1] [1]	[2]
(e) (i)	$12.35 \times 0.01 / 1000 = 1.235 \times 10^{-4}$	[1]	[1]
(ii)	$1.235 \times 10^{-4} \times 1000 / 50 = 2.47 \times 10^{-3}$	[1]	[1]
(iii)	$2.47 \times 10^{-3} \times 64.1 = 0.158327 \text{ g} = 158$ (3 sf only)	[1]	[1]
			[18]
3 (a) (i)	Bond breaking = $\text{Cl-Cl} = 242$ $\text{C-H} = 410 = 652 \text{ kJ}$  Bond forming = $\text{C-Cl} = 340$ $\text{H-Cl} = 431 = 771 \text{ kJ}$  Enthalpy change = $652 - 771 = -119$	[1]  [1]  [1]	[3]
(ii)	UV / High T / sunlight	[1]	[1]

Page 4	Mark Scheme	Syllabus Paper
	Cambridge International AS/A Level – May/June 2015	9701 / 22

Question	Mark Scheme	Mark	Total
(iii)	Initiation $Cl_2 \rightarrow 2Cl\cdot$  Propagation $C_2H_6 + Cl\cdot \rightarrow \cdot C_2H_5 + HCl$ $\cdot C_2H_5 + Cl_2 \rightarrow C_2H_5Cl + Cl\cdot$  Termination $\cdot C_2H_5 + \cdot C_2H_5 \rightarrow C_4H_{10}$  All three names correctly assigned	[1]  [1] [1]  [1]  [1]	[5]
(b) (i)	ethene	[1]	[1]
(ii)	KOH/NaOH  ethanolic <b>AND</b> heat/reflux	[1]  [1]	[2]
(iii)	H <sub>2</sub> <b>AND</b> Pt or Ni (catalyst)	[1]	[1]
			[13]
4 (a) (i)	<b>A</b> = CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CHO <b>B</b> = CH <sub>3</sub> CH <sub>2</sub> CH(CH <sub>3</sub> )CHO <b>C</b> = (CH <sub>3</sub> ) <sub>2</sub> CHCH <sub>2</sub> CHO <b>D</b> = (CH <sub>3</sub> ) <sub>3</sub> CCHO	[1] [1] [1] [1]	[4]
(ii)		[1+1]	[2]
(b) (i)	Fehling's/Benedict's <b>OR</b> Tollens' <b>OR</b> dichromate <b>OR</b> manganate Warm/heat Fehling's/Benedict's =(Brick)-red ppt Tollens' = silver/mirror <b>OR</b> grey/black precipitate Dichromate = orange to green Manganate = purple to colourless <span style="font-size: 2em; vertical-align: middle;">}</span> with the aldehyde/A-D	[1] [1]  [1]	[3]
(ii)	(2,4-)DNP(H)/Brady's reagent  Orange/yellow/red-orange/yellow-orange ppt	[1]  [1]	[2]
			[11]