

CAMBRIDGE INTERNATIONAL EXAMINATIONS

Cambridge International Advanced Level

MARK SCHEME for the May/June 2015 series

9701 CHEMISTRY

9701/51

Paper 5 (Planning, Analysis and Evaluation),
maximum raw mark 30

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2015 series for most Cambridge IGCSE[®], Cambridge International A and AS Level components and some Cambridge O Level components.

© IGCSE is the registered trademark of Cambridge International Examinations.

Page 2	Mark Scheme	Syllabus	Paper
	Cambridge International A Level – May/June 2015	9701	51

Question	Statement	Expected Answer	Mark
1 (a) (i)	M10	$\text{HCOO}^-(\text{aq}) \longrightarrow \text{CO}_2(\text{g}) + \text{H}^+(\text{aq}) + 2\text{e}^-$ $\text{MnO}_4^-(\text{aq}) + 8\text{H}^+(\text{aq}) + 5\text{e}^- \longrightarrow \text{Mn}^{2+}(\text{aq}) + 4\text{H}_2\text{O}(\text{l})$	[1] [1]
(ii)	M6	Magnesium methanoate is $1.312 \text{ mol dm}^{-3}$ $[\text{HCOO}^-(\text{aq})] = 2.624 \text{ mol dm}^{-3}$	[1] [1]
(iii)	M6	Use <u>volumetric apparatus</u> (to measure 5.0 cm^3 / saturated (magnesium) methanoate solution). Make (the above) up to the mark (with water) in a 250 cm^3 volumetric / graduated flask	[1] [1]
(iv)	M3/P4	H^+ is needed for the reaction with manganite Provided the acid is in excess / sufficient / enough, the volume does not matter	[1] [1]
(v)	M5	A pale pink colour	[1]
(vi)	M10	$0.051 \text{ mol dm}^{-3}$	[1]
(vii)	M10	1.28 mol dm^{-3}	[1]

Page 3	Mark Scheme	Syllabus	Paper
	Cambridge International A Level – May/June 2015	9701	51

Question	Statement	Expected Answer	Mark
(b)	P1/P2	(Independent) Temperature (Dependent) Concentration of magnesium methanoate	[1]
(c)	P3	ΔH is positive	[1]
		(An increase in temperature) will favour / promote / increase / a movement in the direction of the endothermic change / reaction	[1]
(d)	P3	Precipitate is formed / barium sulfate is insoluble / insoluble product	[1]
			[15]
2 (a) (i)	D1	$K_c = \frac{[\text{HI}]^2}{[\text{H}_2][\text{I}_2]}$	[1]
(ii)	D1	$K_c = \frac{4y^2}{(a - y)^2}$	[1]

Page 4	Mark Scheme	Syllabus	Paper
	Cambridge International A Level – May/June 2015	9701	51

Question	Statement	Expected Answer	Mark																																				
(b) (i)	D3	<table border="1"> <thead> <tr> <th>a mol dm⁻³</th> <th>a – y mol dm⁻³</th> <th>y mol dm⁻³</th> </tr> </thead> <tbody> <tr><td>0.200</td><td>0.022</td><td>0.178</td></tr> <tr><td>0.500</td><td>0.050</td><td>0.450</td></tr> <tr><td>0.800</td><td>0.252</td><td>0.548</td></tr> <tr><td>1.000</td><td>0.200</td><td>0.800</td></tr> <tr><td>1.500</td><td>0.365</td><td>1.135</td></tr> <tr><td>2.100</td><td>0.570</td><td>1.530</td></tr> <tr><td>2.800</td><td>0.652</td><td>2.148</td></tr> <tr><td>3.400</td><td>0.700</td><td>2.700</td></tr> <tr><td>3.800</td><td>0.867</td><td>2.933</td></tr> <tr><td>4.200</td><td>0.868</td><td>3.332</td></tr> <tr><td>4.900</td><td>1.150</td><td>3.750</td></tr> </tbody> </table>	a mol dm⁻³	a – y mol dm⁻³	y mol dm⁻³	0.200	0.022	0.178	0.500	0.050	0.450	0.800	0.252	0.548	1.000	0.200	0.800	1.500	0.365	1.135	2.100	0.570	1.530	2.800	0.652	2.148	3.400	0.700	2.700	3.800	0.867	2.933	4.200	0.868	3.332	4.900	1.150	3.750	
		a mol dm⁻³	a – y mol dm⁻³	y mol dm⁻³																																			
		0.200	0.022	0.178																																			
		0.500	0.050	0.450																																			
		0.800	0.252	0.548																																			
		1.000	0.200	0.800																																			
		1.500	0.365	1.135																																			
		2.100	0.570	1.530																																			
		2.800	0.652	2.148																																			
		3.400	0.700	2.700																																			
		3.800	0.867	2.933																																			
		4.200	0.868	3.332																																			
		4.900	1.150	3.750																																			
All results for y are to 3 decimal places All values for y are correct																																							
(ii)	D1	All points plotted correctly	[1]																																				
(iii)	E5	Appropriate straight line drawn through the origin	[1]																																				

Page 5	Mark Scheme	Syllabus	Paper
	Cambridge International A Level – May/June 2015	9701	51

Question	Statement	Expected Answer	Mark
(c) (i)	D3/C1	Co-ordinates read correctly from the line	[1]
		Slope of the graph calculated correctly and given to three significant figures with no units.	[1]
(ii)	D3/C1	Uses $\frac{\sqrt{K_c}}{2 + \sqrt{K_c}}$ = gradient (value or y/a) and provides working	[1]
		Gives value of K_c	[1]
(d)	P4	The hydrogen with air / oxygen is explosive at 760K / raised temperature	[1]
(e)	E4	Faster reaction / increased rate	[1]
		The value of K_c would be unaffected	[1]
(f) (i)	E4/C2	The line drawn on the graph has a less steep gradient	[1]
(ii)		The equilibrium constant will be smaller	[1]
			[15]