

| | | |
|---------------|------------------|------|
| Centre Number | Candidate Number | Name |
|---------------|------------------|------|

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS
General Certificate of Education
Advanced Subsidiary Level and Advanced Level

CHEMISTRY

9701/03

Paper 3 Practical Test

October/November 2006

1 hour 15 minutes

Candidates answer on the Question Paper.

Additional Materials: As listed in the Instructions to Supervisors.

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name, including practical session and laboratory where appropriate, in the spaces provided.

Write in dark blue or black pen.

You may use a soft pencil for any diagrams, graphs or rough working.

Do not use staples, paper clips, highlighters, glue or correction fluid.

Answer **all** questions.

You are advised to show all working in calculations.

Use of a Data Booklet is unnecessary.

Qualitative Analysis Notes are provided on pages 7 and 8.

At the end of the examination, fasten your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.


| | |
|---------------------------|--|
| Session | |
| | |
| Laboratory | |
| | |
| For Examiner's Use | |
| 1 | |
| 2 | |
| Total | |

This document consists of **8** printed pages.



- 1 **FA 1** is an aqueous solution of hydrochloric acid, HCl.
FA 2 is aqueous potassium iodate(V) containing 4.93 g dm^{-3} of KIO_3 .
FA 3 is aqueous potassium iodide, KI.
FA 4 is aqueous sodium thiosulphate, $\text{Na}_2\text{S}_2\text{O}_3$.

In the presence of acid, iodate(V) ions react with iodide ions to form iodine and water.



If sodium thiosulphate is present in the reaction mixture the iodine formed is immediately reduced back to iodide and the solution remains colourless.

Iodate(V) ions can therefore be titrated against acid in the presence of an excess of iodide and thiosulphate ions.

You are to use this method to determine the concentration of the hydrochloric acid, **FA 1**.

(a) Dilution of FA 1

Using a burette, run between 34.00 cm^3 and 35.00 cm^3 of **FA 1** into the 250 cm^3 graduated flask, labelled **FA 5**. Record the burette readings in Table 1.1.

Table 1.1 Dilution of FA 1

| | |
|--|--|
| final burette reading / cm^3 | |
| initial burette reading / cm^3 | |
| volume of FA 1 run into the flask / cm^3 | |

Make the solution up to 250 cm^3 with distilled water and **mix thoroughly**.
 Fill a second burette with this diluted acid, **FA 5**.

(b) Titration of IO_3^- with H^+ in FA 5

Pipette 25.0 cm^3 of **FA 2** into a conical flask and use a measuring cylinder to add to the flask 10 cm^3 of **FA 3** and 25 cm^3 of **FA 4**.

Add five drops of bromophenol blue indicator and titrate with **FA 5** until the end-point is reached.

The colour of bromophenol blue changes to yellow at the end-point.

Record your titration results in Table 1.2.

Repeat the titration as many times as you think necessary to obtain accurate results. Make certain that the recorded results show the precision of your practical work.

Table 1.2 Titration of FA 2 with FA 5

| | | | | |
|---|--|--|--|--|
| final burette reading/ cm^3 | | | | |
| initial burette reading/ cm^3 | | | | |
| volume of FA 5 used/ cm^3 | | | | |
| | | | | |

[8]

Summary

25.0 cm^3 of **FA 2** reacted with cm^3 of **FA 5**.

Show which results you used to obtain this volume of **FA 5** by placing a tick (✓) under the readings in Table 1.2.


You are advised to show full working in all parts of the calculations.

- (c) Calculate how many moles of potassium iodate(V), KIO_3 , were pipetted into the conical flask.

[A_r : K, 39.1; I, 127.0; O, 16.0]

[2]

- (d) Calculate how many moles of hydrogen ions, H^+ , reacted with the iodate(V) ions in the flask during the titration.



[1]

- (e) Calculate the concentration, in mol dm^{-3} , of hydrogen ions in **FA 5**.

[1]

- (f) Calculate the concentration, in g dm^{-3} , of hydrochloric acid in **FA 1**.

[A_r : H, 1.0; Cl, 35.5]

[3]

[Total: 15]

- 2 **FA 6** is a mixture of two solids provided in a stoppered boiling tube. One of the solids, **FA 7** is soluble in water; the other, **FA 8** is insoluble in water. Each solid contains one cation and one anion from the ions listed on pages 7 and 8.

In all tests, the reagent should be added gradually with shaking after each addition. Record your observations in the spaces provided.

Your answers should include

- details of colour changes, precipitates formed and the solubility of any precipitate when an excess of the reagent is added,
- details of the test used to identify any gases given off in the reaction.

You should indicate clearly at what stage in a test a change occurs.

Marks are **not** given for chemical equations.

No additional or confirmatory tests for ions present should be attempted.

Candidates are reminded that definite deductions may be made from tests where there appears to be no reaction.

| <i>test</i> | <i>observations [3]</i> |
|---|-------------------------|
| <p>(a) Add water to the boiling-tube labelled FA 6 until it is about half-full. Stopper and shake the tube for 1 minute. Filter the mixture and retain both filtrate and residue for further tests.</p> | |

Tests on the filtrate which contains FA 7

| | |
|---|--|
| <p>(b) To 1 cm depth of the filtrate from (a), in a boiling-tube, add 2 cm depth of aqueous sodium hydroxide.</p> <p>.....</p> <p>Gently warm the solution. Take care as a solution containing sodium hydroxide may 'bump' on heating and eject hot corrosive sodium hydroxide.</p> | |
| <p>(c) To 1 cm depth of the filtrate from (a), in a boiling-tube, add 1 cm depth of aqueous lead(II) nitrate; then</p> <p>.....</p> <p>heat the mixture to boiling point; then</p> <p>.....</p> <p>cool the tube by standing in a beaker of cold water.</p> | |

| <i>test</i> | <i>observations</i> |
|--|---------------------|
| <p>(d) To 1 cm depth of the filtrate from (a), in a test-tube, add 5 drops of aqueous silver nitrate; then</p> <p>.....</p> <p>add 5 cm depth of dilute aqueous ammonia.</p> | |

Use the information in the Qualitative Analysis Notes on pages 7 and 8 to identify the ions present in **FA 7**.

The cation present in **FA 7** is

Which observations indicate the cation you have selected?

.....

.....

..... [1]

The anion present in **FA 7** is

Which observations indicate the anion you have selected?

.....

.....

..... [1]

Tests on the residue which contains FA 8

| <i>test</i> | <i>observations [3]</i> |
|---|-------------------------|
| (e) Transfer the residue to a boiling-tube and add 4 cm depth of dilute hydrochloric acid. Filter the mixture and retain the filtrate. | |
| (f) To 1 cm depth of the filtrate from (e) in a test-tube add, a little at a time, 5 cm depth of aqueous sodium hydroxide. | |
| (g) To 1 cm depth of the filtrate from (e) in a test-tube add, a little at a time, 5 cm depth of dilute aqueous ammonia. | |

Use the information in the Qualitative Analysis Notes on pages 7 and 8 to identify the ions present in **FA 8**.

The cation present in **FA 8** is

Which observations indicate the cation you have selected?

.....

 [1]

The anion present in **FA 8** is

Which observations indicate the anion you have selected?

.....

 [1]

[Total: 10]

QUALITATIVE ANALYSIS NOTES

Key: [ppt. = precipitate]

1 Reactions of aqueous cations

| ion | reaction with | |
|--|---|--|
| | NaOH(aq) | NH ₃ (aq) |
| aluminium, Al ³⁺ (aq) | white ppt. soluble in excess | white ppt. insoluble in excess |
| ammonium, NH ₄ ⁺ (aq) | ammonia produced on heating | |
| barium, Ba ²⁺ (aq) | no ppt. (if reagents are pure) | no ppt. |
| calcium, Ca ²⁺ (aq) | white ppt. with high [Ca ²⁺ (aq)] | no ppt. |
| chromium(III), Cr ³⁺ (aq) | grey-green ppt. soluble in excess giving dark green solution | grey-green ppt. insoluble in excess |
| copper(II), Cu ²⁺ (aq) | pale blue ppt. insoluble in excess | blue ppt. soluble in excess giving dark blue solution |
| iron(II), Fe ²⁺ (aq) | green ppt. insoluble in excess | green ppt. insoluble in excess |
| iron(III), Fe ³⁺ (aq) | red-brown ppt. insoluble in excess | red-brown ppt. insoluble in excess |
| lead(II), Pb ²⁺ (aq) | white ppt. soluble in excess | white ppt. insoluble in excess |
| magnesium, Mg ²⁺ (aq) | white ppt. insoluble in excess | white ppt. insoluble in excess |
| manganese(II), Mn ²⁺ (aq) | off-white ppt. insoluble in excess | off-white ppt. insoluble in excess |
| zinc, Zn ²⁺ (aq) | white ppt. soluble in excess | white ppt. soluble in excess |

[Lead(II) ions can be distinguished from aluminium ions by the insolubility of lead(II) chloride.]

2 Reactions of anions

| <i>ion</i> | <i>reaction</i> |
|---|---|
| carbonate, CO_3^{2-} | CO_2 liberated by dilute acids |
| chromate(VI), $\text{CrO}_4^{2-}(\text{aq})$ | yellow solution turns orange with $\text{H}^+(\text{aq})$; gives yellow ppt. with $\text{Ba}^{2+}(\text{aq})$; gives bright yellow ppt. with $\text{Pb}^{2+}(\text{aq})$ |
| chloride, $\text{Cl}^-(\text{aq})$ | gives white ppt. with $\text{Ag}^+(\text{aq})$ (soluble in $\text{NH}_3(\text{aq})$); gives white ppt. with $\text{Pb}^{2+}(\text{aq})$ |
| bromide, $\text{Br}^-(\text{aq})$ | gives cream ppt. with $\text{Ag}^+(\text{aq})$ (partially soluble in $\text{NH}_3(\text{aq})$); gives white ppt. with $\text{Pb}^{2+}(\text{aq})$ |
| iodide, $\text{I}^-(\text{aq})$ | gives yellow ppt. with $\text{Ag}^+(\text{aq})$ (insoluble in $\text{NH}_3(\text{aq})$); gives yellow ppt. with $\text{Pb}^{2+}(\text{aq})$ |
| nitrate, $\text{NO}_3^-(\text{aq})$ | NH_3 liberated on heating with $\text{OH}^-(\text{aq})$ and <i>Al</i> foil |
| nitrite, $\text{NO}_2^-(\text{aq})$ | NH_3 liberated on heating with $\text{OH}^-(\text{aq})$ and <i>Al</i> foil; NO liberated by dilute acids (colourless $\text{NO} \rightarrow$ (pale) brown NO_2 in air) |
| sulphate, $\text{SO}_4^{2-}(\text{aq})$ | gives white ppt. with $\text{Ba}^{2+}(\text{aq})$ or with $\text{Pb}^{2+}(\text{aq})$ (insoluble in excess dilute strong acid) |
| sulphite, $\text{SO}_3^{2-}(\text{aq})$ | SO_2 liberated with dilute acids; gives white ppt. with $\text{Ba}^{2+}(\text{aq})$ (soluble in excess dilute strong acid) |

3 Tests for gases

| <i>gas</i> | <i>test and test result</i> |
|--------------------------------|--|
| ammonia, NH_3 | turns damp red litmus paper blue |
| carbon dioxide, CO_2 | gives a white ppt. with limewater (ppt. dissolves with excess CO_2) |
| chlorine, Cl_2 | bleaches damp litmus paper |
| hydrogen, H_2 | 'pops' with a lighted splint |
| oxygen, O_2 | relights a glowing splint |
| sulphur dioxide, SO_2 | turns potassium dichromate(VI) (aq) from orange to green |

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.