

**CAMBRIDGE INTERNATIONAL EXAMINATIONS**

Cambridge International Advanced Level

## **MARK SCHEME for the October/November 2015 series**

### **9701 CHEMISTRY**

**9701/43**

Paper 4 (A2 Structured Questions),  
maximum raw mark 100

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the October/November 2015 series for most Cambridge IGCSE<sup>®</sup>, Cambridge International A and AS Level components and some Cambridge O Level components.

© IGCSE is the registered trademark of Cambridge International Examinations.


<b>Page 2</b>	<b>Mark Scheme</b>	<b>Syllabus</b>	<b>Paper</b>
	<b>Cambridge International A Level – October/November 2015</b>	<b>9701</b>	<b>43</b>

Question	Marking Point	Marks	Total Marks
<b>1</b>	<b>(a)</b> ionic bonds break / bonds between $Mg^{2+}$ and $Cl^-$ break  forces / bonds / attractions form between the ions and water	<b>2</b>	
	<b>(b) (i)</b> (the energy change) when 1 mole of a substance dissolves in water / becomes aq	<b>1</b>	
	<b>(ii)</b> $\Delta H_{latt}^{\ominus} MgCl_2 + \Delta H_{sol}^{\ominus} MgCl_2 = \Delta H_{hyd}^{\ominus} Mg^{2+} + 2\Delta H_{hyd}^{\ominus} Cl^-$ $-2524 - 155 = -1925 + 2\Delta H_{hyd}^{\ominus} Cl^-$ $= -377 \text{ kJ mol}^{-1}$	<b>2</b>	
	<b>(iii)</b> magnesium / Mg is higher charge / sodium / Na is smaller charge  magnesium / Mg is smaller / sodium / Na is larger  Mg stronger attraction for water / Na weaker attraction for water any two	<b>2</b>	
<b>(c)</b>	<ul style="list-style-type: none"> <li>solubility decreases</li> <li>lattice energy and hydration enthalpy decrease</li> <li>hydration enthalpy decreases more rapidly / is dominant factor</li> <li>so (enthalpy change of) solution becomes less exothermic / more endothermic</li> </ul>	<b>4</b>	
			<b>[Total: 11]</b>
<b>2</b>	<b>(a)</b> Co $3s^2 3p^6 3d^7 4s^2$ Co <sup>3+</sup> $3s^2 3p^6 3d^6$	[1] [1]	<b>2</b>
	<b>(b) (i)</b> atom or ion, bonded to (one or more), ligands		<b>1</b>
	<b>(ii)</b> any two from: two (or more) oxidation states, catalytic activity, coloured ions or compounds		<b>2</b>

<b>Page 3</b>	<b>Mark Scheme</b>	<b>Syllabus</b>	<b>Paper</b>
	<b>Cambridge International A Level – October/November 2015</b>	<b>9701</b>	<b>43</b>

Question	Marking Point	Marks	Total Marks		
<b>(c)</b>		<b>5</b>			
				<b>transition element species formed</b>	<b>type of reaction</b>
	$\text{Co}^{2+}(\text{aq}) + \text{an excess of NH}_3(\text{aq})$			$[\text{Co}(\text{NH}_3)_6]^{2+}$ or $[\text{Co}(\text{NH}_3)_4]^{2+}$ or $[\text{Co}(\text{NH}_3)_4(\text{H}_2\text{O})_2]^{2+}$	ligand exchange
	$\text{Co}^{2+}(\text{aq}) + \text{OH}^-(\text{aq})$			$\text{Co}(\text{OH})_2$ or $\text{Co}(\text{OH})_2(\text{H}_2\text{O})_4$	precipitation or acid-base
	$\text{Co}^{2+}(\text{aq}) + \text{S}_2\text{O}_8^{2-}(\text{aq})$	$[\text{Co}(\text{H}_2\text{O})_6]^{3+}$ or $\text{Co}^{3+}$ or $\text{Co}_2(\text{SO}_4)_3$	redox or oxidation or reduction of $\text{S}_2\text{O}_8^{2-}$		
<b>(d) (i)</b>	Y 13.4/88.9 or 0.15 Ba 41.2/137 or 0.3 Cu 28.6/63.5 or 0.45 O 16.8/16 or 1	<b>1</b>			
<b>(ii)</b>	= 7/3 or (+) 2.3	<b>1</b>			
<b>(iii)</b>	two Cu are + 2 and one Cu is + 3	<b>1</b>			
			<b>[Total: 13]</b>		



<b>Page 4</b>	<b>Mark Scheme</b>	<b>Syllabus</b>	<b>Paper</b>
	<b>Cambridge International A Level – October/November 2015</b>	<b>9701</b>	<b>43</b>

Question	Marking Point	Marks	Total Marks
3 (a) (i)	<ul style="list-style-type: none"> <li>• <math>\text{Fe}^{2+}</math> and <math>\text{Fe}^{3+}</math> (or suitable compounds),</li> <li>• salt bridge labelled,</li> <li>• one electrode Pt labelled,</li> <li>• one sol<sup>n</sup> 1 mol dm<sup>-3</sup></li> <li>• <math>\text{Cl}^-</math> (or suitable compound),</li> <li>• voltmeter, labelled or V</li> <li>• <math>\text{Cl}_2</math>,</li> <li>• 1 atm or 298K</li> </ul>	2 or 3 marking points = [1] 4 or 5 marking points = [2] 6 or 7 marking points = [3] 8 marking points = [4]	<b>4</b>
	 <p>The diagram shows an electrochemical cell with two half-cells. The left half-cell contains a platinum (Pt) electrode immersed in a solution of iron(II) and iron(III) ions (<math>\text{Fe}^{2+}/\text{Fe}^{3+}</math>). The right half-cell contains a platinum (Pt) electrode immersed in a solution of chloride ions (<math>\text{Cl}^-</math>), with chlorine gas (<math>\text{Cl}_2</math>) being bubbled into the solution from the right. A salt bridge connects the two half-cells. An external circuit connects the two Pt electrodes, containing a voltmeter (V) and two terminals.</p>		
(ii)	$E^\ominus_{\text{cell}} = 1.36 - 0.77 = 0.59 \text{ V}$		<b>1</b>
(b)	yellow/orange/brown		<b>1</b>
(c)	cell voltage increases or becomes more positive $\text{Cl}_2/\text{Cl}^-$ electrode potential increases		<b>2</b>



<b>Page 5</b>	<b>Mark Scheme</b>	<b>Syllabus</b>	<b>Paper</b>
	<b>Cambridge International A Level – October/November 2015</b>	<b>9701</b>	<b>43</b>












Question	Marking Point	Marks	Total Marks
<b>(d) (i)</b>	$\text{H}_2 + 2\text{OH}^- \rightarrow 2\text{H}_2\text{O} + 2\text{e}^-$	<b>2</b>	
	$\text{O}_2 + 2\text{H}_2\text{O} + 4\text{e}^- \rightarrow 4\text{OH}^-$		
	<b>(ii)</b> $2\text{H}_2 + \text{O}_2 \rightarrow 2\text{H}_2\text{O}$		
<b>(iii)</b>	rechargeable/refillable/longer time between charges/longer battery life/less pollution because $\text{H}_2\text{O}$ is the product/ $\text{O}_2$ can be got from the air	<b>1</b>	
			<b>[Total: 12]</b>
<b>4</b>	<b>(a) (i)</b> sketch graph to show a general decrease in m.p	<b>1</b>	
	<b>(ii)</b> giant covalent (C or Si) to metal/metallic (Sn or Pb)	<b>1</b>	
	<b>(b) (i)</b> can react with an acid or base/alkali or can act as an acid or base or has acidic and basic properties	<b>1</b>	
	<b>(ii)</b> $\text{SnO}_2 + 2\text{NaOH} \rightarrow \text{Na}_2\text{SnO}_3 + \text{H}_2\text{O}$ or $\text{SnO}_2 + 2\text{NaOH} + 2\text{H}_2\text{O} \rightarrow \text{Na}_2\text{Sn}(\text{OH})_6$	<b>1</b>	
	<b>(c) (i)</b> $E^\ominus_{\text{cell}} = +1.18$ or $E^\ominus \text{Cr}_2\text{O}_7^{2-}$ greater/more positive than $\text{Sn}^{4+}$ or $E^\ominus (\text{Cr}_2\text{O}_7^{2-}/\text{Cr}^{3+}) + 1.33$ and $E^\ominus (\text{Sn}^{4+}/\text{Sn}^{2+}) + 0.15$	<b>1</b>	
	<b>(ii)</b> $\text{Cr}_2\text{O}_7^{2-} + 3\text{Sn}^{2+} + 14\text{H}^+ \rightarrow 2\text{Cr}^{3+} + 3\text{Sn}^{4+} + 7\text{H}_2\text{O}$ green	<b>2</b>	
	<b>(d) (i)</b> the same substance gets both oxidised and reduced in the reaction or Ge changes oxid. no. +2 to 0 and changes oxid. no. +2 to +4	<b>1</b>	
	<b>(ii)</b> $(\text{CN})_2 + 2\text{NaOH} \rightarrow \text{NaOCN}/\text{NaCNO} + \text{NaCN} + \text{H}_2\text{O}$	<b>1</b>	

Page 6	Mark Scheme	Syllabus	Paper
	Cambridge International A Level – October/November 2015	9701	43

Question	Marking Point	Marks	Total Marks
(iii)	$\begin{array}{c} \text{x} \quad \text{o} \\ \text{N} \equiv \text{C} - \text{C} \equiv \text{N} \\ \text{x} \quad \text{o} \quad \quad \text{o} \quad \text{x} \\ \text{x} \quad \text{o} \quad \quad \text{o} \quad \text{x} \end{array}$	1	
(e) (i)		1	
(ii)	<p>2P<sub>2</sub>: 2 × P≡P = 2 × 489 = 978 kJ mol<sup>-1</sup> <b>and</b>  P<sub>4</sub>: 6 × P–P = 6 × –98 = –1188 kJ mol<sup>-1</sup></p> <p>ΔH = 978 – 1188 = –210 kJ mol<sup>-1</sup></p>	2	
(f) (i)	3NH <sub>4</sub> Cl + 3PCl <sub>5</sub> → 12HCl + P <sub>3</sub> N <sub>3</sub> Cl <sub>6</sub>	1	
(ii)		1	
			<b>[Total: 15]</b>
5 (a) (i)	<p><b>L</b> 2,4-DNPH or Brady's reagent or LiAlH<sub>4</sub> or NaBH<sub>4</sub>  <b>M</b> Fehling's solution or Tollens' reagent or acidified K<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub> or MnO<sub>4</sub><sup>-</sup>  <b>N</b> alkaline I<sub>2</sub></p>	3	
(ii)	CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> CO <sub>2</sub> Na or CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> CO <sub>2</sub> <sup>-</sup> Na <sup>+</sup> or CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> CO <sub>2</sub> H	1	


Page 7	Mark Scheme	Syllabus	Paper
	Cambridge International A Level – October/November 2015	9701	43

Question	Marking Point	Marks	Total Marks
(iii)	yellow precipitate	1	
(iv)	redox or oxidation	1	
(b) (i)	 <p>[1] both curly arrows</p> <p>[1] dipole</p> <p>[1] intermediate</p> <p>two curly arrows [1] dipole [1] intermediate [1]</p>	3	
(ii)		1	
			[Total: 10]

Question	Marking Point	Marks	Total Marks												
6 (a)	<table border="1"> <thead> <tr> <th>reagent</th> <th>organic product</th> <th>non-organic product</th> </tr> </thead> <tbody> <tr> <td>Na</td> <td>  </td> <td>H<sub>2</sub>/hydrogen</td> </tr> <tr> <td>Br<sub>2</sub> (aq)</td> <td>   2 or 3 Br's any position </td> <td>HBr</td> </tr> <tr> <td>CH<sub>3</sub>COCl (l)</td> <td>  </td> <td>HCl</td> </tr> </tbody> </table>	reagent	organic product	non-organic product	Na		H <sub>2</sub> /hydrogen	Br <sub>2</sub> (aq)	 2 or 3 Br's any position	HBr	CH <sub>3</sub> COCl (l)		HCl	4	
	reagent	organic product	non-organic product												
Na		H <sub>2</sub> /hydrogen													
Br <sub>2</sub> (aq)	 2 or 3 Br's any position	HBr													
CH <sub>3</sub> COCl (l)		HCl													
(b) (i)	 <b>E</b>  <b>F</b>	2													




Page 9	Mark Scheme	Syllabus	Paper
	Cambridge International A Level – October/November 2015	9701	43

Question	Marking Point	Marks	Total Marks
(b) (ii)	step 1: $\text{NaNO}_2 + \text{HCl}$ or $\text{HNO}_2$ step 1: $T \leq 10^\circ\text{C}$ step 2: alkaline or $\text{NaOH(aq)}$ or $\text{NaOH}$ solution	3	
			[Total: 9]
7 (a)	<ul style="list-style-type: none"> <li>backbone of sugar-phosphate-sugar-phosphate</li> <li>base bonded to sugar</li> <li>deoxyribose correct label</li> <li>two complementary base pairings e.g A–T or C–G</li> <li>hydrogen bonding/H–bonding between bases, labelled</li> </ul> 	5	
(b)	any <b>two</b> of <ul style="list-style-type: none"> <li>DNA uncoils or unzips</li> <li>hydrogen bonds break or weaken</li> <li>complementary bases join to form a new strand of DNA</li> </ul>	2	



<b>Page 10</b>	<b>Mark Scheme</b>	<b>Syllabus</b>	<b>Paper</b>
	<b>Cambridge International A Level – October/November 2015</b>	<b>9701</b>	<b>43</b>

Question	Marking Point	Marks	Total Marks
<b>(c)</b>	<b>(i)</b> restriction enzymes	<b>1</b>	
	<b>(ii)</b> electrophoresis	<b>1</b>	
	<b>(iii)</b> radioactive substance	<b>1</b>	
	<b>(iv)</b> suspect 3	<b>1</b>	
			<b>[Total: 11]</b>
<b>8</b>	<b>(a) (i)</b> time taken for a compound to travel through the column	<b>1</b>	
	<b>(ii)</b> hydrogen <b>or</b> helium <b>or</b> nitrogen	<b>1</b>	
	<b>(iii)</b> it is more soluble in the stationary phase	<b>1</b>	
	<b>(iv)</b> same functional group <b>or</b> same IMF with stationary phase or same polarity	<b>1</b>	
	<b>(v)</b> % X (= $100 \times 22/76$ ) = <b>29</b> (28.9)	<b>1</b>	
	<b>(b) (i)</b> TMS or tetramethylsilane or Si(CH <sub>3</sub> ) <sub>4</sub>	<b>1</b>	

Page 11	Mark Scheme	Syllabus	Paper
	Cambridge International A Level – October/November 2015	9701	43

Question	Marking Point	Marks	Total Marks																
<b>(ii)</b>	<table border="1"> <thead> <tr> <th>chemical shift <math>\delta</math>/ppm</th> <th>type of proton(s)</th> <th>number of protons</th> <th>splitting pattern</th> </tr> </thead> <tbody> <tr> <td>1.0</td> <td>CH<sub>3</sub>-R</td> <td>3</td> <td>triplet</td> </tr> <tr> <td>2.3</td> <td>CH<sub>2</sub>CO</td> <td>2</td> <td>quartet</td> </tr> <tr> <td>3.7</td> <td>CH<sub>3</sub>O</td> <td>3</td> <td>singlet</td> </tr> </tbody> </table>	chemical shift $\delta$ /ppm	type of proton(s)	number of protons	splitting pattern	1.0	CH <sub>3</sub> -R	3	triplet	2.3	CH <sub>2</sub> CO	2	quartet	3.7	CH <sub>3</sub> O	3	singlet	<b>4</b>	
	chemical shift $\delta$ /ppm	type of proton(s)	number of protons	splitting pattern															
1.0	CH <sub>3</sub> -R	3	triplet																
2.3	CH <sub>2</sub> CO	2	quartet																
3.7	CH <sub>3</sub> O	3	singlet																
<b>(iii)</b>	structure / name of methyl propanoate 	<b>1</b>																	
			<b>[Total: 11]</b>																
<b>9 (a)</b>	C <sub>24</sub> (H <sub>34</sub> )N <sub>2</sub> O <sub>3</sub>	<b>1</b>																	
<b>(b)</b>	ketone    amine    ester	<b>2</b>																	

Page 12	Mark Scheme	Syllabus	Paper
	Cambridge International A Level – October/November 2015	9701	43

Question	Marking Point	Marks	Total Marks
(c) (i)		1	
(ii)		2	
(d)	hydrogen bonding <b>or</b> ion-dipole forces involving lone pair on N atoms, or lone pair on O atoms, or NH <sub>2</sub> groups, or CO <sub>2</sub> groups, or C=O groups, with water	2	<b>[Total: 8]</b>