

**CHEMISTRY****9701/34**

Paper 3 Advanced Practical Skills 2

**October/November 2016**

MARK SCHEME

Maximum Mark: 40

**Published**

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the October/November 2016 series for most Cambridge IGCSE<sup>®</sup>, Cambridge International A and AS Level components and some Cambridge O Level components.

<b>Page 2</b>	<b>Mark Scheme</b>	<b>Syllabus</b>	<b>Paper</b>
	<b>Cambridge International AS/A Level – October/November 2016</b>	<b>9701</b>	<b>34</b>

<b>Question</b>	<b>Answer</b>	<b>Marks</b>
1(a)	<p><b>I</b> Mass of magnesium recorded with /g or (g)  <b>and</b>  initial and final burette readings and volume of hydrogen with unambiguous headings and correct unit.</p> <p>Examiner to calculate 10% and 20% of Supervisor's volume and round this to 1 dp.</p> <p>Candidate's volume compared with Supervisor's volume.</p> <p>Award <b>II</b> if within 20%  Award <b>II</b> and <b>III</b> if within 10%</p>	<p>1</p> <p>1</p> <p><b>3</b></p>
1(b)(i)	<p>Correct calculation moles <math>H_2 = \frac{\text{volume collected}}{24000}</math> to 2 – 4 sf</p> <p>Volume of gas must be correctly calculated.</p>	1
1(b)(ii)	<p>Correctly uses <math>A_r = \frac{\text{mass used}}{\text{(i)}}</math> to 2 – 4 sf</p>	<p>1</p> <p><b>2</b></p>

<b>Page 3</b>	<b>Mark Scheme</b>	<b>Syllabus</b>	<b>Paper</b>
	<b>Cambridge International AS/A Level – October/November 2016</b>	<b>9701</b>	<b>34</b>

<b>Question</b>	<b>Answer</b>	<b>Marks</b>
1(c)(i)	<p>Correct expression</p> $\text{Error in mass} = \frac{0.1 \text{ or } 0.01 \text{ or } 0.001}{\text{mass of Mg}} \times 100^* \text{ (depending on dp of balance)}$ <p>Correct expression</p> $\text{Error in volume} = \frac{0.1 \times 100^*}{\text{volume of gas in (a)}}$	<p>1</p> <p>1</p>
1(c)(ii)	Use a larger mass of magnesium ( <i>for either</i> )/use a balance that reads to more dp ( <i>mass error was larger</i> )/use a burette more precisely calibrated/smaller graduations ( <i>volume error was larger</i> )	<p>1</p> <p><b>3</b></p>



Page 5	Mark Scheme	Syllabus	Paper
	Cambridge International AS/A Level – October/November 2016	9701	34

Question	Answer	Marks
	<p><b>III</b> All accurate burette readings recorded to the nearest 0.05 cm<sup>3</sup>  <i>Do not award this mark if:</i>  <i>50(.00) is used as an initial burette reading;</i>  <i>more than one final burette reading is 50(.00)</i>  <i>any burette reading is &gt; 50(.00)</i></p> <p><b>IV</b> Final uncorrected titre is within 0.10 cm<sup>3</sup> of any previous uncorrected accurate titre.</p> <p>Examiner rounds any accurate burette readings to the nearest 0.05 cm<sup>3</sup>, checks subtractions and then selects the 'best' accurate titres using the hierarchy: identical titres; titres within 0.05 cm<sup>3</sup>; titres within 0.1 cm<sup>3</sup>; etc., to calculate mean correct to 0.01 cm<sup>3</sup>.</p> <p>Examiner compares candidate's titre value with that of the Supervisor.</p> <p><b>V, VI and VII</b>  Award <b>V</b>, <b>VI</b> and <b>VII</b> for <math>\delta \leq 0.20 \text{ cm}^3</math>  Award <b>V</b> and <b>VI</b> for <math>0.20 \text{ cm}^3 &lt; \delta \leq 0.30 \text{ cm}^3</math>  Award <b>V</b> for <math>0.30 \text{ cm}^3 &lt; \delta \leq 0.50 \text{ cm}^3</math></p>	<p>1</p> <p>1</p> <p>1</p> <p>1</p> <p><b>7</b></p>
2(b)	<p>Calculation of mean</p> <p>Check mean titre is correctly calculated from clearly selected values (ticks or working)</p> <ul style="list-style-type: none"> <li>• Candidate must average two (or more) titres where the total spread is <math>\leq 0.20 \text{ cm}^3</math>.</li> <li>• Working must be shown or ticks must be put next to the two (or more) accurate readings selected.</li> <li>• The mean should normally be quoted to 2 dp rounded to the nearest 0.01. [e.g. 26.667 must be rounded to 26.67]</li> </ul> <p>Two special cases where the mean may not be to 2 dp:  allow mean to 3 dp only for 0.025 or 0.075 e.g. 26.325;  allow mean to 1 dp if <b>all</b> accurate burette readings were given to 1 dp (ignoring initial given</p>	1

Page 6	Mark Scheme	Syllabus	Paper
	Cambridge International AS/A Level – October/November 2016	9701	34

Question	Answer	Marks
	<p>as 0) and the mean is exactly correct. [e.g. 26.0 and 26.2 = 26.1 is correct but 26.0 and 26.1 = 26.1 is incorrect.]</p> <p>Do <b>not</b> award this mark if:</p> <ul style="list-style-type: none"> <li>the rough titre was used to calculate the mean;</li> <li>the candidate carried out only 1 accurate titration;</li> <li>burette readings were incorrectly subtracted to obtain any of the accurate titre values;</li> <li><b>all</b> burette readings (resulting in titre values used in the calculation of the mean) are integers.</li> </ul> <p><i>Note: the candidate's mean will sometimes be marked as correct even if it is different from the mean calculated by the examiner for the purpose of assessing accuracy.</i></p>	1
2(c)(i)	<p>Correctly calculates</p> $\frac{2.64}{106 \times 40} = 6.23 / 6.225 / 6.226 \times 10^{-4}$	1
2(c)(ii) and 2(c)(iii)	<p>Correctly uses <b>(i)</b> × 2 <b>and</b> <b>(ii)</b> × 250 <b>(b)</b></p>	1
2(c)(iv)	<p>Correctly calculates</p> $\text{Moles HCl} = \frac{30 \times 1.00}{1000} = 0.03(00)$	1

<b>Page 7</b>	<b>Mark Scheme</b>	<b>Syllabus</b>	<b>Paper</b>
	<b>Cambridge International AS/A Level – October/November 2016</b>	<b>9701</b>	<b>34</b>

<b>Question</b>	<b>Answer</b>	<b>Marks</b>
2(c)(v) and 2(c)(vi)	Correctly uses <b>(iv) – (iii)</b> <b>and</b> $\frac{0.21}{\text{(v)}} \times 2$	1
	Answers to 3 or 4 sf <i>(minimum 4 answers attempted, allow 2 sf in (vi))</i>	1
		<b>5</b>
2(d)(i)	<b>Half</b> the volume needed since 1:1 ratio/1 mole NaOH in equation	1
2(d)(ii)	(Impure) since absorbed/reacted with CO <sub>2</sub> <b>or</b> water vapour/water from the air	1
		<b>2</b>
	<b>Total</b>	<b>15</b>

Page 8	Mark Scheme	Syllabus	Paper
	Cambridge International AS/A Level – October/November 2016	9701	34

Question	Answer	Marks																
	<b>FB 5</b> is CH <sub>3</sub> COCH <sub>3</sub> <b>FB 6</b> is C <sub>2</sub> H <sub>5</sub> OH <b>FB 7</b> is C <sub>2</sub> H <sub>5</sub> CHO <b>FB 8</b> is Cu(NO <sub>3</sub> ) <sub>2</sub>																	
3(a)(i)	<table border="1"> <thead> <tr> <th></th> <th><b>FB 5</b></th> <th><b>FB 6</b></th> <th><b>FB 7</b></th> </tr> </thead> <tbody> <tr> <td>Acidified MnO<sub>4</sub><sup>-</sup></td> <td>no reaction</td> <td colspan="2">Purple to colourless (solution)/(solution) turns colourless</td> </tr> <tr> <td>KI + ClO<sup>-</sup></td> <td colspan="2">(Pale) yellow/cream solid/ppt</td> <td>no reaction</td> </tr> <tr> <td>Tollens'</td> <td colspan="2">no reaction</td> <td>Silver/black/ (dark) grey solid/ ppt/silver mirror</td> </tr> </tbody> </table>		<b>FB 5</b>	<b>FB 6</b>	<b>FB 7</b>	Acidified MnO <sub>4</sub> <sup>-</sup>	no reaction	Purple to colourless (solution)/(solution) turns colourless		KI + ClO <sup>-</sup>	(Pale) yellow/cream solid/ppt		no reaction	Tollens'	no reaction		Silver/black/ (dark) grey solid/ ppt/silver mirror	1  1  1
	<b>FB 5</b>	<b>FB 6</b>	<b>FB 7</b>															
Acidified MnO <sub>4</sub> <sup>-</sup>	no reaction	Purple to colourless (solution)/(solution) turns colourless																
KI + ClO <sup>-</sup>	(Pale) yellow/cream solid/ppt		no reaction															
Tollens'	no reaction		Silver/black/ (dark) grey solid/ ppt/silver mirror															
3(a)(ii)	<b>FB 5</b> is propanone, <b>FB 6</b> is ethanol, <b>FB 7</b> is propanal	1																



Page 9	Mark Scheme	Syllabus	Paper
	Cambridge International AS/A Level – October/November 2016	9701	34

Question	Answer	Marks
3(a)(iii)	Reagent: 2,4–dinitrophenylhydrazine/2,4–DNP(H)/Brady’s reagent Result: propanone and propanal: orange/yellow <b>and</b> solid/ppt (not red) ethanol: no reaction/stays yellow (allow remains colourless) <b>or</b> Reagent: $\text{SOCl}_2/\text{PCl}_3/\text{PCl}_5$ Result: propanone and propanal: no visible reaction/no misty fumes ethanol: steamy/misty fumes (allow white fumes) <b>or</b> Reagent: ethanoic acid + conc $\text{H}_2\text{SO}_4$ (and warm) Result: propanone and propanal: no reaction/ no sweet smell ethanol: sweet/ fruity smell <b>or</b> Reagent: Na Result: propanone and propanal: no reaction/ no bubbles ethanol: effervescence/ bubbling/ fizzing	1 1 <b>or</b> 1 1 <b>or</b> 1 1 <b>or</b> 1 1
3(a)(iv)	Reagent: Fehling’s/Benedict’s/Sandell’s Result: ethanol and propanone: no reaction/stays/turns blue propanal: orange/red/brick-red solid/ ppt	1 1 <b>8</b>
3(b)(i)	(Pale) blue ppt (not dark blue)	1
3(b)(ii)	<b>Black</b> solid	1
3(b)(iii)	Blue/green solution	1

<b>Page 10</b>	<b>Mark Scheme</b>	<b>Syllabus</b>	<b>Paper</b>
	<b>Cambridge International AS/A Level – October/November 2016</b>	<b>9701</b>	<b>34</b>

<b>Question</b>	<b>Answer</b>	<b>Marks</b>
3(b)(iv)	Any <b>two</b> of <ul style="list-style-type: none"> <li>• effervescence/bubbling/fizzing</li> <li>• solid goes pink/brown (allow red-brown)</li> <li>• blue/colour of solution fades (owtte)</li> </ul>	1
3(b)(v)	Oxygen relights glowing splint <b>or</b> nitrogen dioxide is brown	1
3(b)(vi)	$\text{Cu}(\text{NO}_3)_2$	1
3(b)(vii)	$\text{Cu}^{2+}(\text{aq}) + 2\text{OH}^{-}(\text{aq}) \rightarrow \text{Cu}(\text{OH})_2(\text{s})$	1
	<b>Total:</b>	<b>7</b> <b>15</b>